

Algebraic renormalization of antisymmetric tensor matter fields

Vitor Lemes, Ricardo Renan
and
S.P. Sorella

C.B.P.F

*Centro Brasileiro de Pesquisas Fisicas,
Rua Xavier Sigaud 150, 22290-180 Urca
Rio de Janeiro, Brazil*

PACS: 11.15.Bt

REF. 032/94

Abstract

The algebraic renormalization of a recently proposed abelian axial gauge model with antisymmetric tensor matter fields is presented.

1 Introduction

Antisymmetric tensor fields have been introduced since many years [1] and are object of continuous and renewed interests due to their connection with the topological field theories [2]. They are indeed the building blocks of a large class of Schwarz type topological theories [3]. These theories, also known as BF models, can be formulated in any space-time dimension and allow to compute topological invariants which generalize the three dimensional linking number [4]. Moreover they provide an example of ultraviolet finite field theories [5]. Let us point out that the antisymmetric tensor fields of the topological models are gauge fields, i.e. they possess zero modes which have to be gauge fixed in order to have a nondegenerate action.

More recently L. V. Avdeev and M. V. Chizhov [6] achieved the construction of a four dimensional abelian gauge model which includes antisymmetric second rank tensor fields as matter fields rather than gauge fields. The model contains also a coupling of the antisymmetric fields with chiral spinors and a quartic tensor self-interaction term. It exhibits several interesting features among which we underline the asymptotically free ultraviolet behaviour of the abelian gauge interaction. As shown by the authors [6] with an explicit one-loop computation, this is due to the fact that the contribution of the tensor fields to the gauge β -function is negative. Antisymmetric matter fields represent thus a possibility of introducing a new type of interaction in gauge theories.

The aim of this work is to give a regularization independent algebraic analysis of the model. The use of the algebraic method [7] in the present case is motivated by the fact that, besides the presence of γ_5 in the axial couplings of the classical action, the gauge transformations as well as the interaction vertices involving the antisymmetric matter fields explicitly contain the $\varepsilon_{\mu\nu\rho\sigma}$ tensor.

Moreover one has to be sure that the introduction of a new type of matter field does not lead to a new kind of anomaly. Indeed, as

we shall see by exploiting the Wess-Zumino consistency condition [8], the tensor field gives rise to a nontrivial cocycle which is independent from the gauge field and which contains only matter fields. Its numerical coefficient turns out to be related to a set of one-loop Feynman diagrams which are built-up with the interaction vertex between the tensor and the spinor fields. However we will be able to show that the tensor-spinor vertex as well as the matter cocycle can be consistently eliminated by requiring that the model is invariant under an additional discrete symmetry. One is left thus with the usual gauge anomaly of the chiral QED . Furthermore the Adler-Bardeen theorem ensures that the latter is definitively absent if it is absent at the one-loop level, guaranteeing then the ultraviolet renormalizability of the model.

The paper is organized as follows. In Sect. 2 we introduce the classical action and we discuss its stability under radiative corrections. Sect. 3 is devoted to the analysis of the Wess-Zumino consistency condition. In Sect. 4 we present a geometrical derivation of the model in terms of covariant derivatives. Finally, Sect. 5 contains some useful relations and the conventions.

2 The model and its stability

The model is specified by the following classical action which, using the same notation of ref. [6], reads

$$\begin{aligned}
S_{inv} = \int d^4x \Bigg(& -\frac{1}{4h^2} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi}\gamma^\mu\partial_\mu\psi - \bar{\psi}\gamma^\mu\gamma_5 A_\mu\psi + \frac{1}{2}(\partial_\lambda T_{\mu\nu})^2 \\
& -2(\partial^\mu T_{\mu\nu})^2 + 4A_\mu (T^{\mu\nu}\partial^\lambda \tilde{T}_{\lambda\nu} - \tilde{T}^{\mu\nu}\partial^\lambda T_{\lambda\nu}) \\
& +4\left(\frac{1}{2}(A_\lambda T_{\mu\nu})^2 - 2(A^\mu T_{\mu\nu})^2\right) + y\bar{\psi}\sigma_{\mu\nu}T^{\mu\nu}\psi \\
& +\frac{q}{4}\left(\frac{1}{2}(T_{\mu\nu}T^{\mu\nu})^2 - 2T_{\mu\nu}T^{\nu\rho}T_{\rho\lambda}T^{\lambda\mu}\right) \Bigg), \tag{2.1}
\end{aligned}$$

where (h, y, q) are coupling constants and $T_{\mu\nu} = -T_{\nu\mu}$ is a second rank antisymmetric tensor field with

$$\begin{aligned}\widetilde{T}_{\mu\nu} &= \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}T^{\rho\sigma} , \\ \frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}\widetilde{T}^{\rho\sigma} &= -T_{\mu\nu} ,\end{aligned}\tag{2.2}$$

the $\varepsilon_{\mu\nu\rho\sigma}$ tensor being normalized as

$$\varepsilon_{\mu_1\mu_2\mu_3\mu_4}\varepsilon^{\nu_1\nu_2\nu_3\nu_4} = -\delta_{\mu_1}^{[\nu_1}\dots\delta_{\mu_4}^{\nu_4]} .\tag{2.3}$$

We use the Minkowski metric $\eta_{\mu\nu} = \text{diag}(+ - - -)$ and we denote $\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu]$ (see Sect. 5 for the conventions).

Notice that the quadratic term in the antisymmetric field of expression (2.1) is nondegenerate, i.e. $T_{\mu\nu}$ is a matter field. This feature allows also for the introduction of a spinor-tensor interaction (y -term) as well as for a quartic tensor self-interaction (q -term). It is easy to check that the action (2.1) is left invariant by the following gauge transformations [6]:

$$\begin{aligned}\delta A_\mu &= \partial_\mu \omega , \\ \delta \psi &= -i\omega \gamma_5 \psi , \\ \delta \bar{\psi} &= -i\omega \bar{\psi} \gamma_5 , \\ \delta T_{\mu\nu} &= -2\omega \widetilde{T}_{\mu\nu} .\end{aligned}\tag{2.4}$$

Introducing now a covariant Feynman gauge

$$S_{gf} = -\frac{1}{2\alpha} \int d^4x (\partial A)^2 ,\tag{2.5}$$

the gauge-fixed action

$$\Sigma = S_{inv} + S_{gf} ,\tag{2.6}$$

obeys the Ward identity

$$\mathcal{W}(x)\Sigma = -\frac{1}{\alpha}\partial^2\partial A ,\tag{2.7}$$

where $\mathcal{W}(x)$ denotes the local Ward operator¹

$$\mathcal{W}(x) = -\partial_\mu \frac{\delta}{\delta A_\mu} - i \frac{\delta}{\delta \psi} \gamma_5 \psi - i \bar{\psi} \gamma_5 \frac{\delta}{\delta \bar{\psi}} - \widetilde{T}_{\mu\nu} \frac{\delta}{\delta T_{\mu\nu}} .\tag{2.8}$$

¹As usual the spinor derivative $\frac{\delta}{\delta \psi}$ acts from the right to the left.

The classical action Σ is known to be constrained, besides the Ward identity (2.7), by a set of discrete symmetries: i.e. parity \mathcal{P} and charge conjugation \mathcal{C} [9, 10]. They act as

i) Parity \mathcal{P}

$$\begin{aligned} x &\rightarrow x_p = (x^0, -x^i) , & i &= 1, 2, 3 \\ \psi(x_p) &= \gamma^0 \psi(x) , \\ A_0(x_p) &= -A_0(x) , & A_i(x_p) &= A_i(x) , \\ T_{0i}(x_p) &= -T_{0i}(x) , & T_{ij}(x_p) &= T_{ij}(x) . \end{aligned} \tag{2.9}$$

ii) Charge conjugation \mathcal{C}

$$\begin{aligned} \psi &\rightarrow \psi^c = C \bar{\psi}^T , & C &= i\gamma^0 \gamma^2 , \\ A_\mu &\rightarrow A_\mu^c = A_\mu , \\ T_{\mu\nu} &\rightarrow T_{\mu\nu}^c = -T_{\mu\nu} . \end{aligned} \tag{2.10}$$

The fields (A, ψ, T) have respectively dimension $(1, 3/2, 1)$.

In order to study the stability [7] of the model under radiative corrections we look at the most general solution of the equation

$$\mathcal{W}(x) \widetilde{\Sigma} = 0 , \tag{2.11}$$

where $\widetilde{\Sigma}$ is an integrated local polynomial in the fields and their derivatives with dimension four, invariant under parity \mathcal{P} and charge conjugation \mathcal{C} . $\widetilde{\Sigma}$ represents the most general local gauge invariant counterterm which one can freely add at each order of perturbation theory. It can be parametrized as

$$\widetilde{\Sigma} = \widetilde{\Sigma}(A, \psi) + \widetilde{\Sigma}(A, \psi, T) , \tag{2.12}$$

where $\widetilde{\Sigma}(A, \psi)$ depends only on the fields $(A, \psi, \bar{\psi})$ and $\widetilde{\Sigma}(A, \psi, T)$ collects the dependence on the new tensor matter field $T_{\mu\nu}$. The stability condition (2.11), due to the fact that the Ward operator $\mathcal{W}(x)$ is linear, splits into the two conditions

$$\mathcal{W}(x) \widetilde{\Sigma}(A, \psi) = 0 , \tag{2.13}$$

and

$$\mathcal{W}(x)\widetilde{\Sigma}(A, \psi, T) = 0 . \quad (2.14)$$

The first equation (2.13) yields the well known local invariant counterterm of the chiral QED [9]

$$\widetilde{\Sigma}(A, \psi) = \int d^4x \left(-\frac{\rho}{4} F_{\mu\nu} F^{\mu\nu} + \sigma (i\bar{\psi}\gamma^\mu \partial_\mu \psi - \bar{\psi}\gamma^\mu \gamma_5 A_\mu \psi) \right) , \quad (2.15)$$

with (ρ, σ) arbitrary parameters. Turning now to the second term $\widetilde{\Sigma}(A, \psi, T)$ it follows that, using the algebraic identity valid in four dimensions

$$\varepsilon^{\alpha\beta\mu\nu} \partial^\sigma + \varepsilon^{\sigma\alpha\beta\mu} \partial^\nu + \varepsilon^{\nu\sigma\alpha\beta} \partial^\mu + \varepsilon^{\mu\nu\sigma\alpha} \partial^\beta + \varepsilon^{\beta\mu\nu\sigma} \partial^\alpha = 0 , \quad (2.16)$$

it can be parametrized as

$$\begin{aligned} \widetilde{\Sigma}(A, \psi, T) = \int d^4x \bigg(& a(T_{\mu\nu} T^{\mu\nu})^2 + bT_{\mu\nu} T^{\nu\rho} T_{\rho\lambda} T^{\lambda\mu} + c(\partial_\lambda T_{\mu\nu})^2 \\ & + d(\partial^\mu T_{\mu\nu})^2 + e\partial A T_{\mu\nu} \widetilde{T}^{\mu\nu} + mA_\mu T^{\mu\nu} \partial^\lambda \widetilde{T}_{\lambda\nu} \\ & + nA_\mu \widetilde{T}^{\mu\nu} \partial^\lambda T_{\lambda\nu} + p(A^\mu T_{\mu\nu})^2 \\ & + u(A_\mu T_{\lambda\nu})^2 + v\bar{\psi}\sigma_{\mu\nu} T^{\mu\nu} \psi \bigg) , \end{aligned} \quad (2.17)$$

with $(a, b, c, d, e, m, n, p, u, v)$ constant parameters. Condition (2.14) implies that $\widetilde{\Sigma}(A, \psi, T)$ depends only on three parameters, i.e.

$$\begin{aligned} \widetilde{\Sigma}(A, \psi, T) = \int d^4x \bigg(& c\left(\frac{1}{2}(\partial_\lambda T_{\mu\nu})^2 - 2(\partial^\mu T_{\mu\nu})^2 + 4A_\mu T^{\mu\nu} \partial^\lambda \widetilde{T}_{\lambda\nu} \right. \\ & \left. - 4A_\mu \widetilde{T}^{\mu\nu} \partial^\lambda T_{\lambda\nu} + 4\left(\frac{1}{2}(A_\mu T_{\lambda\nu})^2 - 2(A^\mu T_{\mu\nu})^2\right) \right) \\ & + a\left(\frac{1}{2}(T_{\mu\nu} T^{\mu\nu})^2 - 2T_{\mu\nu} T^{\nu\rho} T_{\rho\lambda} T^{\lambda\mu}\right) + v\bar{\psi}\sigma_{\mu\nu} T^{\mu\nu} \psi \bigg) . \end{aligned} \quad (2.18)$$

One sees thus that the most general local gauge invariant counterterm contains five free independent parameters (ρ, σ, c, a, v) . They are easily seen to correspond to a renormalization of the coupling constants (h, y, q) and to field amplitude redefinitions². This proves the stability of the classical action under radiative corrections.

²As it happens in the ordinary QED , the gauge fixing term (2.5) is not renormalized.

3 The Wess-Zumino consistency condition

At the quantum level the action Σ is replaced by a vertex functional

$$\Gamma = \Sigma + O(\hbar) , \quad (3.1)$$

which obeys the broken Ward identity

$$\mathcal{W}(x)\Gamma = -\frac{1}{\alpha}\partial^2\partial A + \mathcal{A}(x) \cdot \Gamma , \quad (3.2)$$

where the insertion $\mathcal{A} \cdot \Gamma$ represents the possible breaking induced by the radiative corrections. According to the Quantum Action Principle [11] the lowest order nonvanishing contribution to the breaking - of order \hbar at least -

$$\mathcal{A} \cdot \Gamma = \mathcal{A} + O(\hbar\mathcal{A}) , \quad (3.3)$$

is a local functional with dimension four, even under charge conjugation \mathcal{C} , odd under parity \mathcal{P} ³ and constrained by the Wess-Zumino consistency condition [8]

$$\mathcal{W}(x)\mathcal{A}(y) - \mathcal{W}(y)\mathcal{A}(x) = 0 . \quad (3.4)$$

The latter stems from the algebraic relation

$$\mathcal{W}(x)\mathcal{W}(y) - \mathcal{W}(y)\mathcal{W}(x) = 0 . \quad (3.5)$$

As it is well known the theory will be anomaly free if condition (3.4) admits only the trivial solution, i.e.

$$\mathcal{A}^{tr}(x) = \mathcal{W}(x)\Delta , \quad (3.6)$$

with Δ an integrated local polynomial with dimension four and even under parity \mathcal{P} and charge conjugation \mathcal{C} . On the other hand nontrivial cocycles of (3.4)

$$\mathcal{A}^{nontr}(x) \neq \mathcal{W}(x)\Delta , \quad (3.7)$$

cannot be reabsorbed as local counterterms and represent real obstructions in order to have an invariant quantum vertex functional.

³This property follows from the fact that the Ward operator $\mathcal{W}(x)$ (2.8) is odd under parity transformations and even under charge conjugation.

To study the Wess-Zumino consistency condition we proceed as before and we write

$$\mathcal{A}(x) = \mathcal{A}_1(A, \psi) + \mathcal{A}_2(A, \psi, T) , \quad (3.8)$$

where $\mathcal{A}_1(A, \psi)$ depends only on $(A, \psi, \bar{\psi})$ and \mathcal{A}_2 contains the field $T_{\mu\nu}$. Equation (3.4) splits thus into the two conditions

$$\mathcal{W}(x)\mathcal{A}_1(y) - \mathcal{W}(y)\mathcal{A}_1(x) = 0 \quad (3.9)$$

and

$$\mathcal{W}(x)\mathcal{A}_2(y) - \mathcal{W}(y)\mathcal{A}_2(x) = 0 . \quad (3.10)$$

The first equation (3.9) yields, modulo trivial cocycles, the usual abelian gauge anomaly [12]

$$\mathcal{A}_1 = r\varepsilon_{\mu\nu\rho\sigma}\partial^\mu A^\nu\partial^\rho A^\sigma . \quad (3.11)$$

Turning to the second equation (3.10) it is not difficult to show the existence of a nontrivial $T_{\mu\nu}$ -dependent cocycle. It reads

$$\mathcal{A}_2 = \eta\bar{\psi}\sigma_{\mu\nu}\gamma_5\psi T^{\mu\nu} , \quad (3.12)$$

with η a numerical coefficient. Notice that expression (3.12) depends only on the matter fields and does not contain the gauge field A . Acting now on the Ward identity (3.2) with the test operator

$$\frac{\delta}{\delta\psi(y)}\frac{\delta}{\delta\bar{\psi}(z)}\frac{\delta}{\delta T_{\mu\nu}(u)} , \quad (3.13)$$

and setting all the fields equal to zero, one easily checks that the numerical coefficient η , at its lowest order, is related to two kinds of one-loop $1PI$ diagrams $\Gamma_{A\psi\bar{\psi}T}^{(1)}$ and $\Gamma_{\psi\bar{\psi}T}^{(1)}$, respectively a set of three box diagrams with four amputated external legs of the type $(A, \psi, \bar{\psi}, T)$ and three triangle diagrams with $(\psi, \bar{\psi}, T)$ as external amputated legs. The appearance of the matter cocycle $\bar{\psi}\sigma_{\mu\nu}\gamma_5\psi T^{\mu\nu}$ could jeopardize the Adler-Bardeen theorem [13, 14] of the gauge anomaly and spoil the renormalizability of the theory.

One is forced then to impose further constraints on the model in order to avoid the presence of the cocycle (3.12). To this purpose let

us remark that all the diagrams which contribute to the coefficient η are built-up by making use of the tensor-spinor three vertex $(\psi\bar{\psi}T)$ of (2.1). Therefore the simplest mechanism which one may impose in order to exclude the matter cocycle is to require from the very beginning that the classical action (2.1) is invariant under an additional discrete symmetry

$$T_{\mu\nu} \rightarrow -T_{\mu\nu} . \quad (3.14)$$

It is apparent that this new invariance forbids the presence in (2.1) of the three vertex $(\psi\bar{\psi}T)$. The latter, as it follows by combining (3.14) with the stability analysis of Sect. 2, will be not reintroduced by the radiative corrections. In addition, requirement (3.14) excludes the appearance of (3.12) as an anomaly in the Ward identity (3.2). We are left then with the gauge anomaly (3.11). However, the Adler-Bardeen theorem [14] ensures that if the coefficient r vanishes at one-loop order it will vanish at all orders of perturbation theory, guaranteeing thus the perturbative renormalizability of the model. As it is well known [9] (see also the detailed discussion of the authors [6]), the vanishing of the coefficient r at the one-loop order in the present abelian case is achieved by introducing a partner for every charged particle with opposite axial gauge charge.

Of course, the discrete symmetry (3.14) will extend to the partner of the tensor matter field $T_{\mu\nu}$.

4 A geometrical construction

In this section we present a simple geometrical derivation of the model. Let us begin by introducing the *covariant derivative*

$$\nabla_\mu T_{\rho\sigma} := (\partial_\mu \widetilde{T}_{\rho\sigma} - 2A_\mu T_{\rho\sigma}) , \quad (4.1)$$

and

$$\nabla_\mu \widetilde{T}_{\rho\sigma} = \frac{1}{2} \varepsilon_{\rho\sigma\alpha\beta} \nabla_\mu T^{\alpha\beta} = -(\partial_\mu T_{\rho\sigma} + 2A_\mu \widetilde{T}_{\rho\sigma}) . \quad (4.2)$$

The name *covariant* is justified by the fact that (∇T) and $(\nabla \widetilde{T})$ transform under the gauge transformations (2.4) as the fields T and \widetilde{T} , i.e.

$$\begin{aligned}\delta(\nabla_\mu T_{\rho\sigma}) &= -2\omega(\nabla_\mu \widetilde{T}_{\rho\sigma}) , \\ \delta(\nabla_\mu \widetilde{T}_{\rho\sigma}) &= 2\omega(\nabla_\mu T_{\rho\sigma}) .\end{aligned}\tag{4.3}$$

This important property allows to construct gauge invariant quantities in a very simple way. It is almost immediate indeed to verify that the term

$$\tilde{S} = \int d^4x \left((\nabla^\mu T_{\mu\nu})(\nabla^\sigma T_\sigma{}^\nu) + (\nabla^\mu \widetilde{T}_{\mu\nu})(\nabla^\sigma \widetilde{T}_\sigma{}^\nu) \right) ,\tag{4.4}$$

is gauge invariant.

Observe that a different contraction of the Lorentz indices

$$\int d^4x \left((\nabla_\mu T_{\rho\nu})(\nabla^\mu T^{\rho\nu}) + (\nabla_\mu \widetilde{T}_{\rho\nu})(\nabla^\mu \widetilde{T}^{\rho\nu}) \right) ,\tag{4.5}$$

identically vanishes due to the property

$$\widetilde{T}_{\rho\sigma} \widetilde{T}^{\rho\sigma} = -T_{\rho\sigma} T^{\rho\sigma} .\tag{4.6}$$

Expression (4.4) is easily computed to be

$$\begin{aligned}\tilde{S} = -\int d^4x \left(\frac{1}{2}(\partial_\lambda T_{\mu\nu})^2 - 2(\partial^\mu T_{\mu\nu})^2 + 4A_\mu T^{\mu\nu} \partial^\lambda \widetilde{T}_{\lambda\nu} \right. \\ \left. - 4A_\mu \widetilde{T}^{\mu\nu} \partial^\lambda T_{\lambda\nu} + 4\left(\frac{1}{2}(A_\mu T_{\lambda\nu})^2 - 2(A^\mu T_{\mu\nu})^2\right) \right) ,\end{aligned}\tag{4.7}$$

i.e. \tilde{S} is nothing but the $(A - T)$ dependent part of the expression (2.1). We have thus recovered in an elegant and more geometrical way the initial invariant classical action S_{inv} . The generalization of the above construction for a nonabelian version of (2.1) is under investigation.

5 Conventions

We give here some useful relations involving the $\varepsilon_{\mu\nu\rho\sigma}$ and the Dirac γ -matrices.

We have

$$\begin{aligned}\widetilde{T}_{\mu\lambda}\widetilde{T}^{\lambda\nu} &= T_{\mu\lambda}T^{\lambda\nu} + \frac{1}{2}\delta_{\mu}^{\nu}T_{\alpha\beta}T^{\alpha\beta} , \\ T_{\mu\lambda}\widetilde{T}^{\lambda\nu} &= \frac{1}{4}\delta_{\mu}^{\nu}T_{\alpha\beta}\widetilde{T}^{\alpha\beta} .\end{aligned}\tag{5.1}$$

From equation (2.16) one gets

$$\begin{aligned}(\partial_{\sigma}\widetilde{T}_{\mu\rho})\widetilde{T}^{\sigma\rho} &= -\frac{1}{2}T^{\alpha\beta}\partial_{\mu}T_{\alpha\beta} - T_{\mu\rho}\partial^{\sigma}T_{\sigma}^{\rho} , \\ \widetilde{T}_{\mu\rho}\partial_{\sigma}\widetilde{T}^{\sigma\rho} &= -\frac{1}{2}T^{\alpha\beta}\partial_{\mu}T_{\alpha\beta} - (\partial_{\sigma}T_{\lambda\mu})T^{\sigma\lambda} ,\end{aligned}\tag{5.2}$$

as well as

$$\widetilde{T}_{\alpha\beta}\partial^2T^{\alpha\beta} = -2T_{\alpha\beta}\partial_{\nu}\partial^{\beta}\widetilde{T}^{\nu\alpha} - 2\widetilde{T}_{\alpha\beta}\partial_{\nu}\partial^{\beta}T^{\nu\alpha} .\tag{5.3}$$

Concerning the γ -matrices, here taken in the Dirac representation, we use [6, 10]:

$$\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 , \quad \gamma_5^2 = 1 ,\tag{5.4}$$

$$\sigma^{\mu\nu} = \frac{i}{2}[\gamma^{\mu}, \gamma^{\nu}] , \quad \frac{i}{2}\varepsilon^{\mu\nu\alpha\beta}\sigma_{\alpha\beta} = \gamma_5\sigma^{\mu\nu} .\tag{5.5}$$

For the charge conjugation matrix C

$$C = i\gamma^0\gamma^2 , \quad C^2 = -1 , \quad C^{-1} = -C ,\tag{5.6}$$

and

$$C\gamma_{\mu}C = \gamma_{\mu}^T , \quad C\gamma_5C = -\gamma_5 = -\gamma_5^T .\tag{5.7}$$

Acknowledgements

We wish to thank José Helayel-Neto for a carefully reading of the manuscript. The *Conselho Nacional de Desenvolvimento Científico e Tecnológico*, *CNPq*-Brazil is gratefully acknowledged for the financial support.

References

- [1] V. I. Ogievetsky and I. V. Polubarinov, *Yad. Fiz.* 4 (1968) 210;
S. Deser, *Phys. Rev.* 178 (1969) 1931;
J. Kalb and P. Ramond, *Phys. Rev.* D9 (1974) 2273;
Y. Nambu, *Phys. Reports* 23 (1976) 250;
- [2] D. Birmingham, M. Blau, M. Rakowski and G. Thompson, *Phys. Reports* 209 (1991) 129;
- [3] A. S. Schwarz, *Baku Intern. Topological Conf. Abstracts*, Vol. 2, (1987) 345;
- [4] G. T. Horowitz, *Comm. Math. Phys.* 125 (1989) 417;
G. T. Horowitz and M. Srednicki, *Comm. Math. Phys.* 130 (1990) 83;
M. Blau and G. Thompson, *Ann. Phys. (N.Y.)* 205 (1991) 130;
- [5] E. Guadagnini, N. Maggiore and S. P. Sorella, *Phys. Lett.* B255 (1991) 65;
C. Lucchesi, O. Piguet and S. P. Sorella, *Nucl. Phys.* B395 (1993) 325;
- [6] L. V. Avdeev and M. V. Chizhov, *Phys. Lett.* B321 (1994) 212;
L. V. Avdeev and M. V. Chizhov, *A queer reduction of degrees of freedom*, preprint *JINR Dubna, hep - th/9407067*;
- [7] C. Becchi, A. Rouet and R. Stora, *Comm. Math. Phys.* 42 (1975) 127;
C. Becchi, A. Rouet and R. Stora *Ann. Phys. (N.Y.)* 98 (1976) 287;
O. Piguet and A. Rouet, *Phys. Reports* 76 (1981) 1;
- [8] J. Wess and B. Zumino, *Phys. Lett.* B49 (1974) 52;
- [9] O. Piguet, “Renormalisation en théorie quantique des champs” and “Renormalisation des théories de jauge”, lectures of the “Troisième cycle de la physique en Suisse Romande” (1982-1983);

- [10] C. Itzykson and J.-B. Zuber, “Quantum field theory”, *McGraw-Hill* 1985;
- [11] Y. M. P. Lam, *Phys. Rev.* D6 (1972) 2145;
Y. M. P. Lam, *Phys. Rev.* D6 (1972) 2161;
T. E. Clark and J. H. Lowenstein, *Nucl. Phys.* B113 (1976) 109;
- [12] S. L. Adler, *Phys. Rev.* 117 (1969) 2426;
J. S. Bell and R. Jackiw, *Nuovo Cim.* 60 (1969) 47;
W. A. Bardeen, *Phys. Rev.* 184 (1969) 1848;
- [13] S. L. Adler and W. Bardeen, *Phys. Rev.* 182 (1969) 1517;
- [14] A. Zee, *Phys. Rev. Lett.* 29 (1972) 1198;
J. H. Lowenstein and B. Schroer, *Phys. Rev.* D7 (1975) 1929;
G. Costa, J. Julve, T. Marinucci and M. Tonin, *Nuovo Cim.* 38A (1977) 373;
G. Bandelloni, C. Becchi, A. Blasi and R. Collina, *Comm. Math. Phys.* 72 (1980) 239;
O. Piguet and S. P. Sorella, *Nucl. Phys.* B381 (1992) 373;
O. Piguet and S. P. Sorella, *Nucl. Phys.* B395 (1993) 661;